Jutge.org

The Virtual Learning Environment for Computer Programming

Induced subgraphs

P12120_en

Novè Concurs de Programació de la UPC - Final (2011-09-21)

Given an undirected graph G = (V, E), any $S \subseteq V$ induces a subgraph G[S] = (S, E'), where E' contains all edges in E that join two vertices in S. Let d(S) denote the minimum degree of the vertices in G[S].

You are given a graph *G* and a size *s*. Which is the maximum degree *d* for which there exists some *S* with at least *s* vertices and such that $d(S) \ge d$?

Input

Input consists of several cases, each with the number of vertices n, the number of edges m, and m pairs $x \ y$ (with $x \neq y$), one for each edge of the graph, followed by s. The vertices are numbered from 0 to n - 1. Assume $1 \le n \le 10^3$, $0 \le m \le n(n - 1)/2$, that there are no repeated edges, and $1 \le s \le n$.

Output

For every case, print the required answer.

S	am	pl	le i	Sample output								
6 0 3	6 1	1	2	2	0	0	3	1	4	2	5	2 1 1
6 0 4	6 1	1	2	2	0	0	3	1	4	2	5	1
2 1 2	1 0											
2 2	0											
3 0 2	2 1	0	2									

Problem information

Author : Salvador Roura Generation : 2024-04-30 15:24:05

© *Jutge.org*, 2006–2024. https://jutge.org