
Similar statements (1)

P70867_en

Catorzè Concurs de Programació de la UPC - Semifinal (2016-06-29)

Consider two infinite horizontal lines A and B , separated ℓ units apart. The line A has m points at the abscissae a_1, \dots, a_m . The line B has n points at the abscissae b_1, \dots, b_n . Given p different indices i_1, \dots, i_p chosen from $\{1 \dots m\}$, and p different indices j_1, \dots, j_p chosen from $\{1 \dots n\}$, define d_k as the Euclidean distance between a_{i_k} and b_{j_k} , that is,

$$d_k = \sqrt{(a_{i_k} - b_{j_k})^2 + \ell^2} .$$

You are given ℓ , p , and the points in A and in B . Pick i_1, \dots, i_p and j_1, \dots, j_p in order to

$$\text{minimize } \sum_{k=1..p} d_k$$

Input

Input consists of several cases, each one with only integer numbers. Every case begins with four strictly positive numbers ℓ , p , m and n . Follow $a_1 \leq a_2 \leq \dots \leq a_{m-1} \leq a_m$. Follow $b_1 \leq b_2 \leq \dots \leq b_{n-1} \leq b_n$. Assume $\ell \leq 10^6$, $p \leq \min(m, n)$, and that the absolute value of each abscissa is at most 10^6 .

Additionally, assume that m and n are at most 1000.

Output

For every case, print the result with four digits after the decimal point. If you use the `long double` type, the input cases have no precision issues.

Sample input

```
1 1 2 2
5 10
9 20

1 2 2 2
5 10
9 20

1000000 4 5 4
300000 300000 300000 300000 300000
-500000 -500000 -500000 -500000

3 2 7 4
0 2 4 6 8 10 12
1 4 7 10
```

Sample output

```
1.4142
14.1730
5122499.3899
6.0000
```

Problem information

Author : Salvador Roura

Generation : 2024-05-02 22:27:49

© *Jutge.org*, 2006–2024.

<https://jutge.org>