
P0012. Diabolical numbers

P79123_en

We say that a number is *diabolical* if it is divisible for the double of the sum of its digits in basis 4. Your task is to write a program that, given a sequence of integers strictly positive finished in -1 , counts how many of them are diabolical.

Your program must include and use the function

```
bool is_diabolical (int n);
```

that indicates if an integer n strictly positive is diabolical or is not.

These are some instances:

n	1	4	6	17	20	23	28	140	255	999999972
n in basis 4	1	10	12	101	110	113	130	2030	3333	323212230213210
sum of the digits	1	1	3	2	2	5	4	5	12	27
diabolical	No	Yes	Yes	No	Yes	No	No	Yes	No	Yes

Input

The input consists of a sequence of integers strictly positive finished in -1 -

Output

Your program must print the quantity of diabolical numbers of the sequence, with six digits. (The inputs will always have less than a million diabolical numbers.)

Sample input 1

-1

Sample output 1

000000

Sample input 2

20 -1

Sample output 2

000001

Sample input 3

17 4 6 20 20 23 140 28 255 999999972 1 2 -1

Sample output 3

000006

Sample input 4

4 4 4 4 4 4 4 4 4 4 4 4 -1

Sample output 4

000012

Problem information

Author : Professorat de P1

Translator : Carlos Molina

Generation : 2013-09-02 15:13:10

© Jutge.org, 2006–2013.

<http://www.jutge.org>