Jutge.org

The Virtual Learning Environment for Computer Programming

Some Hamiltonian paths

Examen parcial d'Algorísmia, FME (2014-11-14)

Consider a directed graph with *n* vertices and all the n(n-1) possible arcs, some of which are painted. How many Hamiltonian paths are in the graph starting at vertex 0, ending at vertex n - 1, and such that they do not traverse two consecutive painted arcs?

Input

Input consists of several cases. Every case begins with *n*, followed by an $n \times n$ matrix, where the position (i, j) has the color of the arc from vertex *i* to vertex *j*. A one indicates a painted arc, and a zero a non-painted arc. The diagonal (which is useless) only has zeroes. You can assume $n \ge 2$.

Output

Sample input	Sample output
2 0 1 1 0	1 1 0
3 0 1 1 1 0 0 1 1 0	4
3 0 1 0 0 0 1 0 0 0	
5 0 1 0 0 0 1 0 1 0 0 0 0 1 1 0 0 1 0 0 1 1 0 0 1 0 0 0 1 0 0 0 0	

For every case, print the number of permutations of the *n* vertices that start at 0, end at n - 1, and do not have three consecutive vertices x, y and z such that the two arcs $x \rightarrow y$ and $y \rightarrow z$ are both painted. The test cases are such that the answer is smaller than 10^6 .

Problem information

Author : Salvador Roura Translator : Salvador Roura Generation : 2024-05-03 08:32:54

© *Jutge.org*, 2006–2024. https://jutge.org