

---

**Disorder Test****X41326\_en**

---

In a strictly increasing ordered sequence of integers  $x_0, \dots, x_r$  (such as 5, 8, 12, 20 for example) each element is strictly larger than the previous one:  $x_i > x_{i-1}$  for all  $i > 0$ , that is,  $x_{i-1} - x_i < 0$ .

A  $k$ -ordered sequence is a sequence where  $x_{i-1} - x_i < k$  for all  $i > 0$ . Thus, a strictly increasing sequence is 0-ordered, and a sequence that is increasing but maybe not strictly (like -3, -1 -1, 4, 4, 7 for example) is 1-ordered.

Larger values of  $k$  represent bounded disorder:  $k$  bounds how smaller than its predecessor each element can be.

Write a function *disorder\_test(k, ls)* that receives a nonnegative integer  $k$  and a list *ls* of integers and checks whether *ls* is  $k$ -ordered, that is, returns `True` if *ls* is  $k$ -ordered, and `False` otherwise.

**Observation**

Only the function will be evaluated. If your submission includes a main program (e.g. with `testmod`), it must be either commented out or inside a condition "`if __name__ == '__main__'`"

**Problem information**

Author: ProAl

Generation: 2026-01-25T15:59:03.783Z

© Jutge.org, 2006–2026.

<https://jutge.org>